ELSEVIER

Forest Ecology and Management 142 (2001) 297–298

Forest Ecology and Management

www.elsevier.com/locate/foreco

Contents of *Forest Ecology and Management*, Volume 142

VOLUME 142, ISSUES 1–3 **1 March 2001**

The physiological basis for the establishment of bare-root larch seedlings
H.M. McKay and J.L. Morgan (Midlothian, UK) 1
Impact of silvicultural system on *Cryphonectria parasitica* incidence and on genetic variability in a chestnut coppice in Central Italy
E. Amorini, M.C. Manetti (Arezzo, Italy), T. Turchetti (Firenze, Italy), A. Sansotta and F. Villani (Porano, Italy) 19
Long-term impacts of logging in a tropical rain forest — a simulation study
A. Huth and T. Ditzer (Kassel, Germany) 33
Individual variability as related to stand structure and soil condition in a Mediterranean oak coppice
F. Logli and R. Joffre (Montpellier, France) 53
The effect of temperature on the radial growth of Scots pine in northernmost Fennoscandia
P. Nöjd (Vantaa, Finland) and P. Hari (Helsinki, Finland) 65
Analysis of diameter–density relationships and self-thinning in non-thinned even-aged Scots pine stands
M. del Río (Palencia, Spain), G. Montero (Madrid, Spain) and F. Bravo (Palencia, Spain) 79
The effects of forest structure and site characteristics on probability of pine twisting rust damage in young Scots pine stands
U. Mattila (Joensuu, Finland), R. Jalkanen and A. Nikula (Rovaniemi, Finland) 89
Impact of grazing on the input of organic matter and nutrients to the soil in a grass-encroached Scots pine forest
A. Smit and A.M. Kooijman (Amsterdam, The Netherlands) 99
Aboveground biomass and nutrient content of five Douglas-fir stands in France
Q. Ponette, J. Ranger, J.-M. Ottorini (Champenoux, France) and E. Ulrich (Fontainebleau, France) 109
The spatial structure of forests in the optimization calculations of forest planning — a landscape ecological perspective
M. Kurttila (Joensuu, Finland) 129
Growth, biomass estimates, and charcoal production of *Acacia drepanolobium* in Laikipia, Kenya
B.D. Okello (Scottsville, South Africa and Nanyuki, Kenya), T.G. O'Connor (Scottsville, South Africa) and T.P. Young (Nanyuki, Kenya and Davis, CA, USA) 143
Estimation of height–diameter curves through multilevel models with special reference to even-aged teak stands
K. Jayaraman (Peechi, India) and J. Lappi (Suonenjoki, Finland) 155
Influence of alternative vegetation management treatments on conifer plantation attributes: abundance, species diversity, and structural diversity
P.M.F. Lindgren and T.P. Sullivan (Vancouver, Canada) 163
Ambrosia and bark beetles (Scolytidae: Coleoptera) in pine and eucalypt stands in southern Brazil
C.A.H. Flechtmann (Ilha Solteira, Brazil), A.L.T. Ottati (Botucatu, Brazil) and C.W. Berisford (Athens, GA, USA) 183
A height increment equation for young ponderosa pine plantations using precipitation and soil factors
F.C.C. Uzoh (Redding, CA, USA) 193
Effects of fire on bark beetle presence on Jeffrey pine in the Lake Tahoe Basin
T. Bradley and P. Tueller (Reno, NV, USA) 205
Multivariate point processes and spatial variation of species diversity
K. Shimatani (East Lansing, MI, USA) 215
Above- and below-ground production of trees and other vegetation on contrasting aspects in western Montana: a case study
D.W. Coble (Nacogdoches, TX, USA), K.S. Milner (Missoula, MT, USA) and J.D. Marshall (Moscow, ID, USA) 231

Attempting restoration of wet tropical forests in Costa Rica
A. Carl Leopold (Ithaca, NY, USA), R. Andrus (Binghamton, NY, USA), A. Finkeldey (Lansing, NY, USA) and D. Knowles (Greenville, NC, USA) 243
Biotic integrity of *Platanus wrightii* riparian forests in Arizona: first approximation
J.C. Stromberg (Tempe, AZ, USA) 251
Bird responses to partial retention timber harvesting in central interior British Columbia
A.N. Lance and M. Phinney (Prince George, BC, Canada) 267
Dry weight estimation of the aboveground components of *Melaleuca quinquenervia* trees in southern Florida
M.B. Rayachhetry, T.K. Van, T.D. Center (Fort Lauderdale, FL, USA) and F. Laroche (West Palm Beach, FL, USA) 281
Time trajectories of mass and density in a *Chamaecyparis obtusa* seedling population
K. Ogawa (Nagoya, Japan) 291

Contents of *Forest Ecology and Management*, Volume 142 297